

SMART
eNAT®

Smart Delivery System for Nucleic Acids Preservation Medium





Copan patented SMART push & turn activation and delivery system is a post-collection, high-performance yet intuitive cap that avoids unwanted media spillage or contact, facilitating self-sampling at home. SMART-eNAT® combines the SMART-cap™ with eNAT® and FLOQSwabs®, stabilizing and preserving RNA/DNA for prolonged periods. SMART-eNAT® is intended to collect, transport, and preserve clinical specimens to be analyzed by nucleic acids amplification techniques.



SMART-CAP™

Unique push & turn activation and delivery system that makes sample collection easy, while guarding against unwanted spillage or contact.



EASY TO USE

SMART-eNAT® has been designed keeping the end-user in mind. Only a few simple steps are required to collect the sample.



VERSATILE

SMART-eNAT® is available paired with three different swab shafts specific for vaginal, buccal, and rectal collection or in a kit with a swab for microbiome research purposes.



eNAT®

A liquid-based molecular medium that quickly inactivates viruses and bacteria while preserving nucleic acids at room temperature to reduce turnaround time and increase safety.



Preservation

eNAT® Performance

Copan eNAT® medium preserves nucleic acids for:

- Up to 4 weeks at RT and 4°C¹
- Up to 6 months at -20°C to -80°C

According to the vast scientific literature, eNAT® characteristics successfully preserved nucleic acids from fecal samples up to 30 days at room temperature².

FLOQSwabs®

Cut out for everyone

FLOQSwabs® ensure a quick, capillarity-driven sample uptake and a superior elution of the biological specimen.

Its combination with SMART-eNAT® expands your downstream diagnostic testing possibilities, by offering an unmatched specimen collection in many anatomical sites.

Discover why we call them “the perfect collection device.”



Fields of application

Prenalytics made different



Respiratory Diseases^{3,4,5}



Gastrointestinal Diseases^{6,7,8}



STI & HPV^{9,10,11}



Genetics & Microbiome^{12,13,14,15}

Combinations

The SMARTer way!

SMART-eNAT® expresses its full potential when paired with the right FLOQSwabs®.

Choose the vaginal swab, rectal swab or the buccal hDNAfree FLOQSwabs®.




1 ml eNAT® medium in the SMART-cap®



1 regular FLOQSwabs® for Vaginal Sample

or



1 regular FLOQSwabs® for Rectal Sample

or



1 regular FLOQSwabs® hDNAfree for Buccal Sample



Laboratory

Handling & processing

Samples collected with Copan SMART-eNAT® are suitable for commercial nucleic acids extraction and amplification platforms.

Scientific literature reports sample collection and transport with eNAT® prior to many downstream processes:

- Molecular-based assays ^{9,15,16,17,18,19,20,21}
- Next Generation Sequencing ^{13,14,22,23,24}

Self-Collection

Collect your nucleic acids like a pro

Reaching end-users and donors in the comfort of their homes is critical to delivering highly personalized health information and suggestions. That's why the Self-collection of biological samples has become increasingly important in genetics, genomics, and microbiome research. **SMART-eNAT® enables a clean and safe collection of nucleic acids at home, making DNA and RNA sampling easy and accessible to many users in many different settings.**



Ordering information

SMART-eNAT® is available in combination with FLOQSwabs®.

Cat. N.	Description	Pack size	Sample*
6E099S	1 ml eNAT® medium in the smart cap 	300 pieces	
5U175S01	1 regular FLOQSwabs® 	600 pieces	Vaginal
5U173S01	1 regular FLOQSwabs® hDNAfree Free 	600 pieces	Rectal
5U174DS01	1 regular FLOQSwabs® 	600 pieces	Buccal
Cat. N.	Description	Pack size	Sample*
70U001N**	1 ml eNAT® medium in the smart cap + 1 regular FLOQSwabs®  	300 pieces (6 boxes of 50 pieces)	Nasal, throat, vaginal, skin, rectal, wound, buccal and faeces

*Suggested table. Please refer to your GLP procedures to choose the most appropriate device for the specific sampling site
 ** NO CE IVD - not for clinical use

Scientific references

All the studies we cited in this product focus are listed here.

1. Hasan MR et al. (2012) Short-term stability of pathogen-specific nucleic acid targets in clinical samples. *J Clin Microbiol.* 50(12):4147-50
2. Young RR et al. (2020) Long-term stability of microbiome diversity and composition in fecal samples stored in eNAT medium. *Microbiologyopen.* 9(7):e1046
3. Lee J et al. (2018) Evaluation of Allplex Respiratory Panel 1/2/3 Multiplex Real-Time PCR Assays for the Detection of Respiratory Viruses with Influenza A Virus subtyping. *Ann Lab Med* 38(1): 46-50
4. Ordonez-Mena JM et al. (2020) Relationship between microbiology of throat swab and clinical course among primary care patients with acute cough: a prospective cohort study. *Fam Pract* 37(3):332-339
5. Mhimbira F et al. (2019) Prevalence and clinical significance of respiratory viruses and bacteria detected in tuberculosis patients compared to household contact controls in Tanzania: a cohort study. *Clin Microbiol Infect* 25(1): 107.e1-107.e7
6. Liu J et al. (2016) Optimization of Quantitative PCR Methods for Enteropathogen Detection. *PLoS One* 11(6): e0158199
7. Pernica J et al. (2018) Short-course antimicrobial therapy for paediatric respiratory infections (SAFER): study protocol for a randomized controlled trial. *Trials* 19(1):83
8. Pichon M et al. (2020) Diagnostic Accuracy of a Noninvasive Test for Detection of *Helicobacter pylori* and Resistance to Clarithromycin in Stool by the Amplidag H. *pylori*+ClariR Real-Time PCR Assay. *J Clin Microbiol* 58(4): e01787-19
9. Yar DD et al. (2016) Genotypic characterisation of human papillomavirus infections among persons living with HIV infection; a case-control study in Kumasi, Ghana. *Trop Med Int Health* 21(2):275-82
10. Juliana NCA et al. (2020) The Prevalence of Chlamydia trachomatis and Three Other Non-Viral Sexually Transmitted Infections among Pregnant Women in Pemba Island Tanzania. *Pathogens* 9(8):625
11. Wendland EM et al. (2019) Evaluating sexual health in sex workers and men who have sex with men: the SMESH cross-sectional protocol study. *BMJ Open.* 9(11):e031358
12. Susic D et al. (2020) Microbiome Understanding in Maternity Study (MUMS), an Australian prospective longitudinal cohort study of maternal and infant microbiota: study protocol. *BMJ Open* 10: e040189
13. Rooney AM et al. (2020) Each Additional Day of Antibiotics Is Associated With Lower Gut Anaerobes in Neonatal Intensive Care Unit Patients. *Clin Infect Dis* 70(12): 2553-2560
14. Ata B et al. (2019) The Endobiota Study: Comparison of Vaginal, Cervical and Gut Microbiota Between Women with Stage 3/4 Endometriosis and Healthy Controls. *Sci Rep* 9, 2204
15. Roy C et al. (2020) Performance Evaluation of the Novodiag Bacterial GE+ Multiplex PCR Assay. *J Clin Microbiol.* 58(10):e01033-20
16. Thomas PPM et al. (2019) Sexually Transmitted Infections and Behavioral Determinants of Sexual and Reproductive Health in the Allahabad District (India) Based on Data from the ChlamIndia Study. *Microorganisms.* 7(11):557
17. Falaschi Z et al. (2020) Chest CT accuracy in diagnosing COVID-19 during the peak of the Italian epidemic: A retrospective correlation with RT-PCR testing and analysis of discordant cases. *Eur J Radiol.* 130:109192
18. Welch SR et al. (2020) Analysis of Inactivation of SARS-CoV-2 by Specimen Transport Media, Nucleic Acid Extraction Reagents, Detergents, and Fixatives. *J Clin Microbiol.* 58(11):e01713-20
19. Narchi H et al. (2020) Nasopharyngeal Isolates from a Cohort of Medical Students with or without Pharyngitis. *Sultan Qaboos Univ Med J.* 20(3):e287-e29
20. Nagel A et al. (2020) Characterization of a universal screening approach for congenital CMV infection based on a highly-sensitive, quantitative, multiplex real-time PCR assay. *PLoS One.* 15(1):e0227143
21. Lee YM et al. (2020) Monitoring environmental contamination caused by SARS-CoV-2 in a healthcare facility by using adenosine triphosphate testing. *Am J Infect Control* 48(10): 1280-1281
22. van Pamel J et al. (2020) Alterations of Gut Microbiota and the Brain-Immune-Intestine Axis in Patients With Relapsing-Remitting Multiple Sclerosis After Treatment With Oral Cladribine: Protocol for a Prospective Observational Study. *JMIR Res Protoc.* 9(7):e16162
23. Ciardiello T et al. (2020) Effects of Fermented Oils on Alpha-Biodiversity and Relative Abundance of Cheek Resident Skin Microbiota. *Cosmetics* 7(2):34
24. Cieplik F et al. (2020) Oral Health, Oral Microbiota, and Incidence of Stroke-Associated Pneumonia-A Prospective Observational Study. *Front Neurol.* 11:528056



This document may contain product information otherwise not accessible or valid in your country. Please be aware that Copan Italia S.p.A. does take any responsibility for accessing such information which may not comply with any valid legal process, regulation, registration or usage in the country of your origin. Product clearance and availability restrictions may apply in some Countries. Please refer to Copan website (www.copangroup.com) to view and/or download the most recent version of the brochure. This document is mainly intended for marketing purposes, always consult product insert for complete information. The use of this product in association with diagnostic kits or instrumentation should be internally validated by the user. ©2023 Copan Italia. All rights reserved. The trademarks mentioned herein are property of Copan Italia S.p.A.
Code: JMKPF013Ro3.EN



@copangroup

Copan Italia s.p.a.
Via Francesco Perotti 10,
25125 Brescia, Italy

t | f +030 2687211
@ | info@copangroup.com
www.copangroup.com